



*10th International Conference on
Pattern Recognition and
Information Processing*

PRIP'2009

Proceedings

19-21 May, 2009

Minsk, Belarus

УДК 681.3

PATTERN RECOGNITION AND INFORMATION PROCESSING: Proceedings of the Tenth International Conference (19 - 21 May, Minsk, Republic of Belarus), - Minsk, 2009. – 400p.

Proceedings contain papers accepted for publication and presentation at the Tenth International Conference “Pattern Recognition and Information Processing” that will be held in May 19-21, Minsk, Belarus. Proceedings are prepared for publication by PRIP’2009 Program Committee and Belarusian Association for Image Analysis and Recognition.

Proceedings will be useful for students and researchers working in the following areas: pattern recognition and image analysis; knowledge processing and representation; knowledge based decision support system, fuzzy mathematics and systems, applications of pattern recognition and image analysis.

Papers are approved by Program Committee and are printed as submitted by authors.

Сборник включает доклады десятой Международной конференции “Распознавание образов и обработка информации”, которая состоится 19-21 мая 2009 года в Минске (Беларусь). Подготовлен к выпуску программным комитетом конференции PRIP’2009 и Белорусской ассоциацией по анализу и распознаванию изображений.

Доклады конференции будут полезны студентам и специалистам в следующих областях: распознавание образов и анализ изображений, обработка и представление знаний, системы поддержки принятия решений, основанные на знаниях, нечеткая математика и системы, приложения распознавания образов и анализа изображений.

Статьи одобрены Программным комитетом и печатаются в виде, представленном авторами.

Editors:

Victor Krasnoproshin, Sergey Ablameyko, Rauf Sadykhov

ISBN 985-6453-90-9
ISBN 985-64589-5

© Belarusian State University (Belarus),
2009

Organized by

BELARUSIAN STATE UNIVERSITY, *Faculty of Applied Mathematics and Computer Sciences*

In cooperation with

- Belarusian State University of Informatics and Radioelectronics
- United Institute of Informatics Problems of the National Academy of Sciences of Belarus
- Belarusian Association for Image Analysis and Recognition
- FE Invention Machine Belarus Ltd.
- Scientific & Technological Association “INFOPARK”

In collaboration with

- Belarusian Republican Foundation for Fundamental Research
- FE Byelex Multimedia Products Ltd.
- International Associations for Pattern Recognition



InventionMachine

PRIP'2009 COMMITTEES

Chairman of the Conference

Krasnoproshin V. Belarus

Vice-Chairmen of the Conference

Ablameyko S. Belarus

Sadykhov R. Belarus

International Scientific Committee

G. Borgefors	Sweden	M. Petrou	UK
D. Chetverikov	Hungary	G. Sanniti di Baja	Italy
T. Deserno	Germany	V. Shmerko	Canada
V. Donskoy	Ukraine	I. Sovpel	Belarus
J. Gil Aluha	Spain	N. Trush	Belarus
I. Gurevich	Russia	A. Tuzikov	Belarus
R. Kasturi	USA	S. Uchida	Japan
Yu. Kharin	Belarus	V. Viergever	Netherlands
B.C. Lovell	Australia	E. Zaitseva	Slovakia
P. Mandrik	Belarus	A. Zhilinskas	Lietuva
A. Marcelli	Italy	Yu. Zhuravlev	Russia
E. Oczeretko	Poland		

Local Organizing Committee

V. Obraztsov - chairman

A. Bobkov

S. Gafurov

A. Dobrovolskij

V. Konakh

T. Korbut

N. Rubashko

A. Valvachov

Welcome Words from the PRIP'09 Chairman



On behalf of the Organizers it is my great pleasure to welcome you at the 10th International Conference Pattern Recognition and Information Processing (PRIP). PRIP Conference is the main biennial event of the Belarusian Association for Image Analysis and Recognition. The PRIP'09 is being held at the Belarusian State University - in one of the leading and oldest Belarusian universities, from May 19-21, 2009. PRIP is held in cooperation with other scientific establishments. PRIP'09, as the previous Conferences is endorsed by the International Association for Pattern Recognition (IAPR). It is also sponsored by some other organizations.

It is my pleasure to inform that 126 papers have been submitted for PRIP'09 from 30 countries (256 authors). All submitted papers have been reviewed by Program Committee members together with referees. As a result we have selected 88 papers for inclusion in the PRIP'09 scientific program.

Such a large-scale Conference can not take place without the help of many people in the Conference organization. I would like to express my sincere gratitude to all members of the Program Committee and referees who carefully reviewed and selected the papers, to the Organizing Committee who has been working hard for the organization of the Conference and preparation of PRIP'09 Proceedings book, to all authors for the high scientific quality of their papers. Special thanks to our organization - Belarusian State University. Many thanks to everyone who took part in preparation of the Conference.

I am happy to welcome you to Minsk and hope that you will have good impressions and memories of PRIP'09 when you will open this book.

Victor Krasnoproshin

PRIP'09 Chairman
Minsk, Belarus, April 2009

Image Processing for Morphological Investigation of the Carcinoma Thyroid Gland: Modern Facilities

Nedzved A.¹⁾, Fridman M.²⁾, Belotserkovsky A.¹⁾

1) United Institute of Informatics Problems, Minsk, Belarus nedzveda@newman.bas-net.by

2) Republican Centre for Thyroid Tumours, Minsk, Belarus

Abstract: This paper describes automation methods for diagnostics of the thyroids gland carcinoma. It based on image processing technology and calculation of characteristics from image. Methods are used for investigation of histological samples.

Keywords: Medical image processing, calculation of characteristics.

1. INTRODUCTION

Digital image processing became one of mainstems in computer science that bring technical progress in many fields of human activity. An application of methods to be discussed here is intended for efficiency and reliability increasing of accepted decisions since histological diagnostics of oncological diseases is one of urgent problems in medical practice. It can be solved by using of modern computer system of the image analysis [1-5].

Morphological analysis of histological samples is usually carried out by analysis of images made with different optical magnifications (low and high, for instance). In the first case architectonics is investigated under magnification about 50-100X. In the second case cells and endocellular structures are investigated at magnification of 400-1000X in compliance with class of cells.

Simple thresholding methods based on brightness histogram analysis are applied to define patterns of tissue. These patterns are optionally modified then by operations of mathematical morphology. While it should consist of the integral objects with approximately repeating borders of cells, it preserve a hierarchy of objects (nucleus and nucleolus). After morphometrical analysis the formalized characteristics are calculated, that create a basis for allocations of conclusion.

2. MORPHOLOGICAL CHARACTERISTICS AT LOW OPTICAL MAGNIFICATION

There are fragments of tissue represented on image at low optical magnification of specimen (fig. 1). Cells are to be detected as small contrast objects [2], so after segmentation image contains binary patterns which include both nucleus and noise (fig. 2). Then, full histology functional representation (fig. 3) is brought by distance map transformation [2, 7]. The described method is used for investigation of tumour's morphometric characteristics of thyroid glands.

It is considered that at low magnification any new tumour formation may be investigated by following characteristics:

- blood circulation infringement;
- adjournment of salts of calcium;
- texture of colloids;
- characterization of a fibrous tissue distribution and growth features;

- the architectonic of tumours;
- degree of expressiveness of productive inflammation (mononuclear infiltration);
- size of follicles;
- characteristic of papillae;
- presence of capsule.




Fig.1 – Initial histological image of carcinoma of the thyroid gland, x100




Fig.2 – The binary image of cells, x100

The formalization of these characteristics is carried out in two ways: definition of specific objects and calculation of morphometric characteristics.

Such features as fibrosis and/or capsules, infringement of blood circulation, suspension of salts of calcium can be defined by specialized segmentation operations. For example, fibrosis is characterized by tissue that is optically homogeneous. For its definition it is possible to use threshold segmentation and morphological filtration. This allows to modify borders and remove geometrical

noise.



Fig.3 – Segmented binary image of tumour tissue

There are three kinds of fibrosis (fig. 4) and each of them is characterized by certain geometrical complexity.




Fig.4 – Character of tumour fibrosis

The character of tumour growth depends on regions of fibrosis (fig. 5). It is characterized by properties of a capsule. In such a case the capsule should have fibrosis-like optical properties, since in the ideal case the capsule's border is narrow and closed. Therefore image processing by the morphological filtration demands satisfaction of additional conditions to control over geometrical characteristics of the defined objects.




Fig.5 –Features of tumour growing

The colloid is described as "liquid", "dense", "pseudo-

calcification". So, optical density is used for its characterization and can be calculated as follows:

$$OD_{x,y} = -\log \left(\frac{I_{x,y} - I_{\min}}{I_{\max} - I_{\min}} \right), \quad (1)$$

where OD_{xy} – optical density; I_{xy} – brightness of a point; I_{\min} and I_{\max} – minimal and maximal brightness correspondingly.

A tumor's architectonics is presented by follicular, solid and papillary structures (fig. 6).




Fig.6 – Variants of structural atypia in tumours and tumor-like diseases of the thyroid

The expressiveness degree of immune inflammation is formalized by the area of particles and distance distribution between nucleuses.

For the description of follicles (small, large and average) the area of objects is used, but characterization of papillary structures is more difficult. For papillary thyroid cancer "true" papillae (with fibrosis and prominent vasculature in stalk), papillae's hyperplasia (with follicles in similar pouching – in goiter) are usually under consideration. Formalization of these characteristics requires additional investigation, but it is possible to use the characteristic of convex for the simplified case, which must not exceed the value of 0,64 for papillary hyperplasia.

3. MORPHOLOGICAL CHARACTERISTICS AT HIGH OPTICAL MAGNIFICATION

Internuclear distance plays a leading role in the complex analysis. It may be found from pixel value on distance map (fig. 7). Value of each point from the distance map corresponds to the nearest distance to closest nucleus.




Fig.7 –Analysis of internuclear characteristics a) the tissue image, b) a distance map of internuclear region.

Internuclear distances between cells and shapes of regions have to be defined for each type tumours. For example, describe the shape of papillary thyroid cancer

the convexity feature is used (fig. 8):

$$convexity = \frac{convex\ perimeter}{perimeter} \quad (2)$$

If object is convex, this value is equal to 1 while less than 1 if hollows are exist.

The similar description is used for density investigation. It is called "solidity" and defined as relation of the common and convex areas [2]:

$$solidity = \frac{AREA}{convex\ area} \quad (3)$$

If the object has no holes and hollows, this size is equal to 1, if else it has less value.



Fig.8 – Objects of different convexity and solidity: convexity and solidity are equal 1; convexity is equal 0,483, solidity - 0,782; convexity is equal 0,349, solidity - 0,592.

At high optical magnification characteristics of cells and their structures become basics for description. Therefore nucleus and cytoplasm are objects of interest in this case. For example, cytoplasm is described as amphophilic or weak-basophilic, eosinophilic and water-clear. Formalization of dimension (fig. 9) is simple and corresponds to the description of optical density.




Fig.9 – The cytoplasm features

Changes of nucleus at various diseases of a thyroid gland are divided on three groups (fig. 10):

- 1) intranuclear changes:
 - a) an inclusion;
 - b) a groove;
 - c) presence of a large nucleolus;
 - d) a mitosis;
- 2) chromatin:
 - a) regular distributed;
 - b) basophilic clumpy;
 - c) highly dispersed;
- 3) form of nucleus:
 - a) ovoid;
 - b) round;
 - c) bizarre;
- 4) sizes of a cell;
- 5) presence of symplaste.




Fig.10– The colour image of cell with a nuclear

Pathognomonic changes are found out in a nuclear (fig. 11) and are presented by cytoplasmic inclusions and grooves.




Fig.11– The colour image of cell

Grooves look like narrow extensive objects crossing a nuclear from one edge to another (fig. 12). Inclusions are characterized by vial-like objects. Object's contents are painted the same as cytoplasm (fig. 12).




Fig.12 – Pathological changes of a nuclear: a) groove, b) mitosis, chromosome backlog in body, c) a mitosis, backlog of a fragment of a chromosome in a metaphase, d) inclusions.

Nuclear analysis brings topological properties of objects which allow to define pathognomonic structure type for a papillary thyroid cancer:

- groove – if there are more than 2 points of crossings with nuclear border ;
- mitosis – if there are more than one large object;
- mitosis – if camber of object is less 0,8;
- inclusion – if the brightness of object is high.

To study a nuclear structure it is suitable to use hierarchical image of its. Such an image can be obtained as a result of segmentation process. At the first stage the Otsu thresholding is performed (fig. 13) to binarize an image which has been obtained by difference of orthogonal components in LAB color coordinate system [7].




Fig.13 – Threshold segmentation of objects on the histological image: a) the original image, b) the result o binarisation

Morphological opening operation divides then the merged objects to find topological characteristics. To connect broken lines of nucleus borders closing morphological operation must be carried out after. Finally filling operation of the closed contours finishes the process (fig. 14a). Geometrical noises are removed by the analysis of the sizes and the shape. For example, for identification of nuclear structure initial binary image is used. At this step subtraction of borders (fig. 14) define intranuclear objects. The result of such operation is "hierarchical" multiphase image (fig. 14d).




Fig.14 – Binary image of a nucleus (a), grooves (b), nucleolus (c) and hierarchical image of a nucleus (d)

The idea of hierarchy of areas is easily realized with help of computer facilities

Table 1. Values of pixels for different elements in structure of cages

Object	Operation	Separate value	Values after association
Nuclear	$1 \ll 0$	1	1
Grooves or inclusions	$1 \ll 1$	2	3
Nucleolus	$1 \ll 2$	4	7

It is possible to define about eight levels of hierarchy. Advantage of the method described here lies in one hierarchical image. It allows to describe all levels of

classification. As a result this image allows to obtain full information about nuclear.

Thus, if there are large objects, (furrows or inclusions) inside nucleus, the value of pixel corresponds to 3. For nucleolus or chromatin the pixel equal to 7. At the similar organization it is possible to operate easily with various nuclear components. It is possible to investigate topological structure of a nuclear by such association of bits in the hierarchical image. The nuclear characteristics are easy formalized for the chromatin description by using of such concepts as geometrical density of objects, intercept and the sum of chords.

4. NUMERICAL CHARACTERISTICS

Developed methods and algorithms have been tested on images of histological samples of the thyroid gland diseases: adenoma (15 cases), goiter (12) and papillary thyroid cancer (21). Following dependences have been received:

Table 2. Mean values of cell's nucleus

disease	area, 10^{-6}	Max. diameter, 10^{-6}	Min. diameter, 10^{-6}	Optical density	Optical range
adenoma	26.7	7.21	4.749	0.316	0.039
goiter	22.4	6.09	4.659	0.56	0.066
cancer	38.2	7.97	6.067	0.29	0.048
norm	21.0	6.68	4.11	0.47	0.057

Table 3. Dispersion of values of cell's nucleus

disease	area, 10^{-6}	Max. diameter, 10^{-6}	Min. diameter, 10^{-6}
adenoma	121.10	2.00	1.66
goiter	87.63	1.38	1.05
cancer	138.98	2.05	1.49
norm	24.68	1.18	0.62

Table 4. Characteristics of follicles and intercellular space

disease	Mean diameter, 10^{-6}	Diameter dispersion, 10^{-6}
adenoma	146.6	3403.00
goiter	127.3	2847.38
cancer	42.5	421.05
norm	83.7	1606.3

Thus average values of nuclear area for a papillary thyroid cancer are much bigger than for thyroid cells in norm or at benign conditions. The dispersion of nuclear area for a papillary cancer is more than other. Therefore an extreme variety of nuclear changes for carcinomas are exists in this case. It is necessary to notice smaller diameter of follicles in a follicular variant of a papillary cancer in comparison with a crawl and an adenoma. That feature can give additional criterion of differential diagnostics.

5. CONCLUSION

Histological characteristics are formalized by using of methods of the images analysis and calculations of characteristics for raster objects. The similar approach allows to automate histological research. It gains good results for quality and speed of diagnostics. The greatest problems are caused by process of structure definition of nucleus and cells, because these images have small contrast and no accurate borders. But described methods

allow constructing full functional of histology investigation without dependence on complexity of initial images by the expense of application of binarisation and segmentations methods.

6. ACKNOWLEDGEMENT

This work is partially supported by ISTC projects #B-1636 and #B-1682.

7. REFERENCES

- [1] Ablameyko S. and Belotserkovsky A. Medical Image Segmentation Using Homogeneity and Color Characteristics Histograms. *Vesti of National Academy of Sciences of Belarus*, Minsk, Belarus, №2, 2003, pp. 76-83.
- [2] S.Ablameyko, A.Nedzved. *Processing of optical images of cellular structures in medicine*. UIIP NAS Belarus, Minsk. 2005. p.155.
- [3] S.Ablameyko, D.Lagunovsky et al. From Cell Image Segmentation to Differential Diagnosis of Thyroid Cancer *Proceedings of 16th Intern. Conf. on Pattern Recognition*, Quebec, Canada, 11-15 August 2002, vol. 1, pp. 763-766.
- [4] N.Lassouaoui, L.Hamami, A.Zerguerras Segmentation and classification of biological cell images by a multifractal approach *International Journal of Intelligent 18* (6), (2003), p. 657-678.
- [5] A.Nedzved, A.Belotserkovsky, S.Ablameyko Computer systems of histology image analysis in Belarus. *Annual Proceeding of Medical Science*, Bialystok, Poland: MUB, 50 (2005). pp.23-26.
- [6] P.K.Sahoo, S.Soltani, A.K.C.Wong, Y.C.Chen, A survey of thresholding techniques. *Computer Vision, Graphics, and Image processing*, 41 (1988). pp. 233-260.
- [7] T.Tanaka, T.Joke., T.Oka Cell nucleus segmentation of skin tumour using image processing *Proceeding of the 23 int.conf. on EMBS*, 2001, pp.2716-2719.